

Drosophila immune cell migration and adhesion during embryonic development and larval immune responses

Aparna Ratheesh, Vera Belyaeva and Daria E Siekhaus



The majority of immune cells in *Drosophila melanogaster* are plasmatocytes; they carry out similar functions to vertebrate macrophages, influencing development as well as protecting against infection and cancer. Plasmatocytes, sometimes referred to with the broader term of hemocytes, migrate widely during embryonic development and cycle in the larvae between sessile and circulating positions. Here we discuss the similarities of plasmatocyte developmental migration and its functions to that of vertebrate macrophages, considering the recent controversy regarding the functions of *Drosophila* PDGF/VEGF related ligands. We also examine recent findings on the significance of adhesion for plasmatocyte migration in the embryo, as well as proliferation, trans-differentiation, and tumor responses in the larva. We spotlight parallels throughout to vertebrate immune responses.

Address

IST Austria, Am Campus 1, 3400 Klosterneuburg, Austria

Corresponding author: Siekhaus, Daria E (daria.siekhaus@ist.ac.at)**Current Opinion in Cell Biology** 2015, **36**:71–79This review comes from a themed issue on **Cell adhesion and migration**Edited by **Michael Sixt** and **Erez Raz**<http://dx.doi.org/10.1016/j.ceb.2015.07.003>0955-0674/© 2015 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Immune cells are essential for survival, as they eliminate both foreign invaders and endogenous pathologies [1,2]. While vertebrates utilize a complex set of innate and adaptive immune cells, *Drosophila melanogaster* relies on an innate immune system consisting of only three cell types, jointly called hemocytes, to play a broad range of roles [3]. Plasmatocytes, the functional equivalent of vertebrate macrophages, are 95% of all *Drosophila* immune cells prior to infection and will be the focus of this review. They influence development [4*,5,6] and physiology [7] as well as defend against bacteria [8,9], fungi [8], viruses [10], and cancer [11,12**]. Plasmatocytes migrate actively during embryonic development [13] and pupation [14], as well as during responses to wounds [15,16]. In the larva, many of their positions are due to regulated adhesion [17**,18]. We have sought to avoid overlap with


two recent excellent reviews [9,19]; here we focus on the conservation between *Drosophila* plasmatocyte and vertebrate macrophage migration in embryos, and examine the adhesion involved in larval plasmatocyte physiology and tumor responses. We highlight questions throughout that we consider intriguing for further exploration.

Conservation of embryonic macrophage migration paths and functions in *Drosophila* and vertebrates

Much of the embryonic migration of *Drosophila* plasmatocytes occurs along paths where their function is required for further development. Plasmatocytes are specified in the anterior mesoderm in the ventral side of the head [20,21]; they then ingress [22] and split into three main routes, two of which have at this time been shown to have clear developmental relevance (Figure 1). In route 1, plasmatocytes move over the yolk sac to the tip of the germband (route 1a). They then invade the epithelia of the extended germband [13,23,24*] on their way to kidney-like organs called the renal tubules (route 1b); plasmatocytes secrete collagen IV which facilitates BMP signaling required for the proper positioning of these organs [4*]. These plasmatocytes then migrate along the posterior ventral nerve cord (vnc) (1c), eventually joining the cells moving from their birthplace towards the posterior along the vnc in route 2; all along the vnc, plasmatocytes engulf apoptotic midline glia and facilitate vnc condensation [5,6,25]. Route 3 along the forming heart [16] has not yet been shown to have a developmental role but in any case serves to further disperse plasmatocytes in preparation for larval immune functions.

These embryonic migration paths and their purposes show similarities with those of vertebrate macrophages formed during primitive hematopoiesis (see Table 1) [26–28]. As in *Drosophila*, macrophages in zebrafish are specified in the anterior ventral mesoderm. They then move onto the yolk sac as in route 1 [29]; this step also precedes their penetration of epithelial tissues [30], and phagocytosis of apoptotic cells of the nervous system [31]. The precursors of mouse macrophages are also born in the anterior mesoderm and move onto the yolk sac; there they form blood islands in which they mature [32] before appearing in the head [33] and seeding the brain where they develop into microglia [34]. Movement analogous to route 2 along the vnc is observed in zebrafish and the chick, in which macrophages move into the spinal cord from anterior to posterior after their population of the head [29,35]. Mouse macrophages infiltrate the

Figure 1



Plasmatocyte migration routes and their functional roles during embryonic development. Schematic of two embryos (early Stage 12 on the left and Stage 14 on the right) showing that plasmatocytes specified in the head mesoderm migrate along three main routes during embryonic development. One sub population migrates in Stage 12 over the yolk sac to the edge of the extended germband indicated by an asterisk (route 1a). They then penetrate the germband epithelium and cluster around the renal tubules where they secrete collagen IV which ensheathes the tubules (route 1b). These and other plasmatocytes that have entered the germband continue along the posterior ventral nerve cord (vnc, route 1c in left embryo, route 1 in right embryo). Another subpopulation migrates out from the head (route 2 in both embryos) along the anterior ventral nerve cord. In both of these routes plasmatocytes engulf apoptotic midline glia. The third group of plasmatocytes migrates along the developing heart also towards the posterior of the embryo (route 3 in both embryos). Arrows indicate the migration routes.

developing kidney interstitium and may stimulate growth and ureteric bud branching [36]. Postnatally mouse macrophages also facilitate the branching of the mammary gland, a process requiring Bone morphogenetic protein (BMP) signalling [37,38]. Macrophage remodeling, although not secretion, of collagen appears to be involved [39]. Thus macrophages influence development in both *Drosophila* and vertebrates and migrate developmentally to many of the same tissues. This routing helps populate different vertebrate tissues with the resident macrophages that play later essential physiological and immunological roles [40].

PDGF/VEGF ligands in *Drosophila* and vertebrate macrophage migration

PDGF/VEGF-related ligands (PvFs) have been thought to mediate migration along all three embryonic routes in *Drosophila* but this idea is now contested. The original hypothesis rested on the findings that each path expresses one of the 3 PvFs [13,16] and that loss of function of the ligands or their plasmatocyte expressed receptor, the PDGF/VEGF-related Receptor, PVR, causes defects in movement along each route [13,16,23,41]. However, interpretation of these experiments is complicated; PVR signaling is also required for plasmatocyte survival [23]. PVR activation of Mbc and Rac has been implicated in its migratory function in another cell type [42,43], and signaling through Akt/Tor, and MEK/ERK in its role in hemocyte survival [13,23,42,44,45]. Thus to definitively

demonstrate a migratory role for these ligands or their receptor requires the migration defects caused by their absence to remain when cell survival is restored. This has been shown for PVR and Pvf2/3 in penetration of the germband in route 1 [23,41]. In route 2 the importance of PVR [16] is established but that of PvFs is not yet clear. One lab showed strong migratory defects after RNAi of Pvf2 and 3, but did not assess effects on plasmatocyte survival [16]. Another rescued survival and restored the migratory defects seen in a deletion affecting the two PvFs, however this deletion causes only a reduction, not the elimination, of Pvf2 expression [41]. A role in route 3 is likely as migration there fails in the absence of only one Pvf [16]; eliminating two is required to see strong survival defects [13,23]. Whether these PvFs are acting as chemoattractants is another open question. When Pvf2 is over-expressed in areas the plasmatocytes normally cross, it triggers plasmatocyte accumulation, which could be caused by attraction or adhesion [13,16,25]. PvFs have not been used to redirect plasmatocytes to a new area, as was demonstrated with another migratory cell type, border cells [46]. Expression of Pvf2 or a dominant active (DA) form of PVR in the plasmatocytes themselves should block migration if a chemotactic response is required for guidance. Each appeared not to, but the expression was turned on only after much migration had already commenced [41] and in a background in which the endogenous protein was still present, albeit for Pvf2 at reduced levels. Thus the potential migratory functions for

Table 1**Summary of *Drosophila* plasmatocyte embryonic migration routes, factors, functions and conservation with those of vertebrate macrophages**

<i>Drosophila</i> plasmatocyte route	<i>Drosophila</i> route description	<i>Drosophila</i> ligands and receptors	<i>Drosophila</i> experiments and caveats	<i>Drosophila</i> functional relevance	Vertebrate route conservation	Vertebrate receptors involved	Vertebrate functional conservation
1a	Over the yolk sac to edge of posterior germband	PVR independent	PVR null mutant still moves up to edge of germband [13].	None yet identified.	Zebrafish and mouse macrophage precursors move over yolk sac [29,32,33].	VEGFR-2 needed for macrophage precursors to move onto yolk sac blood islands in mouse [47,48].	
1b	Penetration between posterior germ band epithelia on the way to the renal tubules.	PVR Pvf2 (Pvf3)	PVR null mutant rescued for cell survival shows no movement into germ band [23]. Pvf2/3Δ shows no movement into germ band. Phenotype rescued just by Pvf2 expression [41].	Collagen IV secretion to facilitate BMP signaling needed for renal tubule development [4].	Kidney infiltration by macrophages seen in mouse [36]. Epithelial penetration seen in zebrafish [30].	CSF1R needed for epithelial penetration in zebrafish (Fig. 9E,F in [30]).	Remodeling of collagen involved in mammary gland development seen in mouse [39].
1c	Along the posterior ventral nerve cord (vnc)	PVR Pvf2&3?	Pvf2 and 3 RNAi knockdown show migration defects along vnc, cell survival not assessed [16]. Pvf2/3Δ mutant defects restored upon rescue of cell survival [41]. Yet Δ mutant is not a complete null: reduces Pvf2, truncates Pvf3.	Engulfment of apoptotic midline glia [5,6], vnc condensation.	Zebrafish macrophages appear in posterior nerve cord (Fig. 8S in [29]).		Apoptotic neural cells engulfed in zebrafish [31].
2	Along the anterior vnc	PVR Pvf2&3?	PVR null mutant rescued for cell survival shows little movement along anterior vnc [16]. Pvf experiments and caveats same as above [41].	Engulfment of apoptotic midline glia [5,6], vnc condensation.	Zebrafish (Fig. 9E,F, in [30]), chick.	CSF1R (Fig. 9E,F in [30,35]).	Apoptotic neural cells engulfed in zebrafish [31].
3	Along the forming heart	PVR Pvf2	PVR null mutant rescued for cell survival shows little movement along forming heart [16]. Pvf2 transposon insert mutant and RNAi showed defects [16]. Cell survival not assessed but lacking one Pvf does not cause strong survival defects [13,23].	None yet identified.			

Each row corresponds to a route taken by *Drosophila* plasmatocytes during their embryonic migration. For each route, successive columns indicate the signals and receptors currently known to be required for the indicated migration and then the experiments underlying that conclusion and their caveats. A question mark indicates that the corresponding molecule has been contradictorily identified both as a plasmatocyte migratory cue and as solely a survival factor, as discussed in the caveat column. Further columns illustrate the potential conservation of the *Drosophila* plasmatocyte routes with those of vertebrate macrophages and the vertebrate receptor required for the vertebrate route indicated. The final column delineates the potential conservation of a functional role with vertebrates.

Pvfs are to facilitate invasion in route 1, mediate adhesion or guidance on several routes, or all of the above.

Even if the Pvfs do guide migration, many questions remain. Movement along the first step (1a) of route 1 up to the germband can occur even in the absence of PVR [13], implying the existence of another migratory cue for this step. Each of the three main routes that the plasmatocytes split into contains Pvfs [13,16], thus how the cells decide which path to follow is unclear. Finally, along all three paths, consecutive waves of plasmatocytes move towards one source of Pvf, but then move beyond it to another. Thus, if Pvf guides movement during normal development, mechanisms must exist within the migrating hemocyte streams to create a gradient from the successive concentrations of Pvf expression, as in the zebrafish lateral line [47,48]. Alternatively, contact with the leading hemocyte could induce tissues to downregulate Pvfs or upregulate sequestering receptors [49] so that the leading hemocyte would receive greater signal from targets further ahead. This would require, however, that subsequent hemocytes follow cues not from their surroundings but from other hemocytes.

The closest vertebrate orthologs of *Drosophila* PVF are Vascular Endothelial Growth Factor (VEGF) and Platelet derived growth factor (PDGF). These can guide the migration of macrophages during development and of monocytes, the precursors of macrophages, during physiological responses. VEGF Receptor 2 (VEGFR-2) is needed for macrophage precursors to appear in blood islands in mice [50]; this is thought to be due to a defect in their migration as VEGFR-2-mutant cells can differentiate properly *in vitro* [51]. A role for PDGFR β in migration of macrophage precursors to blood islands or from the yolk sac has not been assessed, but it is not required for the developmental migration of hematopoietic stem cells from the fetal liver [52]. Purified VEGF can guide human monocytes across endothelial monolayers [53]; both VEGF and PDGF can direct monocyte chemotaxis *in vitro* [54–56]. The next closest ortholog of *Drosophila* PVR, after PDGFR and VEGFR, is the Colony Stimulating Factor 1 Receptor (CSF1R), which is involved in monocyte/macrophage precursor chemotaxis [57]. Interestingly, in zebrafish the invasion of macrophages from the yolk sac into the brain, retina and epidermis depends on CSF1R, which starts to be expressed in pre-macrophages maturing in the yolk sac [30]. Thus as evolution proceeded, the migratory functions of *Drosophila* PVR may have been split between VEGFR, PDGFR, and CSF1R [58] during development and immunological responses.


Modulation of adhesion during the *Drosophila* plasmatocyte life cycle

Integrin adhesion plays an essential and dynamic role in facilitating and influencing the migration of

plasmatocytes in the embryo. Integrin affinity is regulated by the GTPase Rap1 [59], as in vertebrates in which both of these proteins are required for the movement of neutrophils and monocytes between endothelial cells out of the vasculature [60]. *Drosophila* plasmatocytes also penetrate a tissue barrier as they move into the germband along route 1b and analogously require α -Integrin, Inflated, as well as Dizzy, a GEF for Rap1 [24*] for this step. Modulation of this adhesion appears to be crucial as the GTPase RhoL, which regulates Rap1 localization and thus Integrin affinity, is essential for this process. Plasmatocytes could use Integrins to bind the germband's epithelial cells and change their junctional properties to permit penetration, as vertebrate monocytes do while exiting blood vessels [61]. Alternatively, Integrins could facilitate homotypic adhesion since plasmatocytes migrate in chains during germband entry, contacting the rear of the cell ahead [24*]; indeed strong plasmatocyte β -Integrin dependent clustering can be induced at later stages by over expressing Dizzy or Rap1DA [59]. In contrast, at these later stages, overlap that arises normally between lamellipods leads to repulsion, facilitating the dispersal and movement of hemocytes [62*,63**]. The contacting lamellipods form an adhesion that leads to the coordinated reorganization of the colliding cytoskeletal networks and a build up of accumulated tension [63**]; its release seems to propel repulsion. Integrins could be involved in this event, as in its absence the cells maintain contact longer and move more slowly away from one another [64]. Thus plasmatocytes seek contact at early stages and are repelled by it at later ones; this change could be due to a temporal shift in plasmatocyte signaling pathways downstream of Integrins.

Embryonic plasmatocytes persist into the larval stage, but in this period active migration plays a more limited role than adhesion. During all larval stages, plasmatocytes circulate passively in the lymph that bathes the internal organs and are then recruited to tissue surfaces and wound sites by adhesion [65,66]. In the early larvae, plasmatocytes also home based on cues provided by neurons to segmentally repeated pockets between muscles and the epidermis where they attach to the internal surface of the body wall [17**,67] (Figure 2). Localization in these pockets permits these sessile plasmatocytes to undergo a faster rate of division, receive survival signals, and trans-differentiate. Their presence at these locations requires Eater, a hemocyte specific EGF-like repeat receptor [68*]. These sites maintain their attractive capacities over time because plasmatocytes return after mechanical disruption displaces them [17**]. Yet this localization is also dynamic; at later larval stages these plasmatocytes undergo exchanges between the body wall pockets [17**]. Trans-differentiation of a few plasmatocytes into crystal cells occurs in a Notch-dependent manner even in the

Figure 2



Larval hemocytes exist in sessile patches and in circulation. **(a)** Schematic showing hemocyte distribution in a 3rd instar larva. Hemocytes colonize segmentally repeated epidermal-muscular pockets found along the side of the embryo (indicated in one segment by the red box) and attach to the internal body wall from early larval stages. At later stages hemocytes are also found in association with the dorsal vessel (indicated in one segment by a purple bracket). Sessile hemocytes undergo exchanges between the pockets on the body wall (shown with bi-directional arrows) and during immune challenges return to circulation. Cartoons depicted below correspond to the boxed region in the larva and demonstrate different sessile hemocyte behaviors. **(b)** Sessile hemocytes in the epidermal-muscular pockets cluster around the oenocytes and associate with cells of the peripheral nervous system (PNS), which are essential for their trophic survival. Hemocyte association with the sessile compartment requires the plasmatocyte specific EGF-like repeat receptor, Eater. Hemocytes also exchange between sessile patches and the circulation. **(c)** Plasmatocytes attached to the sessile patches undergo proliferation. **(d)** Plasmatocytes attached to the sessile compartment can trans-differentiate into crystal cells in a Notch dependent manner.

absence of the wounds and parasites that the crystal cells serve to melanize [69–71]. These crystal cells remain in the pockets as long as plasmatocytes express Eater and are also located there [68^{*}]. Immune challenge leads to the return to circulation of plasmatocytes and crystal cells [72,73]; if the infecting agent is a parasite, these released sessile plasmatocytes also transdifferentiate into lamellocytes which wrap around the invaders [72].


What molecular mechanisms trigger the alterations in adhesion underlying their dynamic cycling between pockets in the normal larva or their mobilization in the infected one is an open question. Expression in plasmatocytes of two genes, either of which should disrupt Wg signaling, releases sessile plasmatocytes [74]. Constitutive Toll signaling in the fat body can also lead to disruption of the plasmatocyte pattern [73]. These results argue that unknown external signals heralding the presence of infection can decrease adhesive strength directly in sessile hemocytes or in the muscles or epidermis they

bind to. As Eater also binds to bacteria to permit their phagocytosis [75,76], plasmatocytes that are triggered to leave and then encounter bacteria might be temporarily precluded from rebinding to the pockets. Whether plasmatocytes returning after exposure to pathogens can shift the proliferation or differentiation rate of the sessile ones and thus act analogously to macrophages and dendritic cells presenting antigen to T cells in lymph nodes is an intriguing area to explore [77]. In any case, larval plasmatocyte adhesion in these pockets is required for their expansion and responses to infection, behaviors also observed in vertebrate tissue resident macrophages which they have been proposed to be analogous to [27,33,78,79].

Plasmatocyte tumor responses initiated by adhesion

Circulating plasmatocytes are captured by adhesion to larval tumors where they can block or promote aberrant cell growth, depending on the tumor type (Figure 3). Tumors induced in salivary glands solely by oncogenic

Figure 3



Tumor associated hemocytes can lead to tumor promotion and invasion or tumor regression. Schematic depiction of *Drosophila* hemocyte and tumor interactions. Plasmacytes are recruited to adhere to tumors of all genetic types. The further responses of both cell types depend on the genetic makeup of the tumor, as indicated below. In tumors induced in salivary glands by Ras^{V12}, lamellocytes and crystal cells are recruited to the tumor, leading to its encapsulation. In tumors induced in imaginal discs by mutations in the polarity genes, *scribble* and/or *discs large*, plasmacyte derived Eiger causes tumor cells to upregulate Pvf1, leading to further plasmacyte proliferation. Plasmacyte Eiger also triggers tumor inhibition in combination with factors from the fat body. Eiger is a transmembrane protein; it may act through direct contact with tumor cells or be secreted after cleavage. In imaginal disc tumors deficient for *scribble* but overexpressing Ras^{V12}, plasmacyte derived Eiger mediates a switch in tumor response from *in situ* residence to invasive migration.

Ras^{V12} are bound by plasmacytes, lamellocytes, and crystal cells. These immune cells encapsulate and melanize the transformed tissues, isolating it as they do with wasp eggs [80]. Tumors elicited in imaginal discs by mutations in the polarity genes, *scribble*, *discs large* or *lethal giant larvae* [81], lead to the adhesion of plasmacytes at areas where the basement membrane is disrupted [11]. These plasmacytes inhibit tumor growth by producing Eiger, the only identified member in *Drosophila* of the Tumor Necrosis Factor (TNF) α superfamily [11,12**]. Plasmacyte Eiger leads to a positive feedback loop of tumor control; it induces tumor cells to die and to express Pvf1 which results in plasmacyte proliferation through PVR signalling [12**].

Finally, if the tumors induced by polarity gene mutations in imaginal discs also express Ras^{V12}, plasmacytes are again captured from the circulation by adhesion, but lead to a different response. Eiger produced by these plasmacytes causes not tumor death, but rather overgrowth and invasive migration [82,83*]. This final case shows similarities to vertebrates, in which tumor associated macrophages promote tumor functions through TNF α as well as pro inflammatory cytokines [84]. There are likely to be common signals, perhaps a disrupted basement membrane, through which all *Drosophila* tumors induce plasmacyte adhesion. Yet there must also be distinct tumor signaling pathways that lead to the specific plasmacyte responses to different

tumor types and divergent tumor responses to plasmatocyte produced Eiger.

Conclusions

Due to the relative ease of genetic manipulation and imaging in *Drosophila*, its immune system serves as an excellent system to study how cellular migration occurs within diverse *in vivo* environments. While migration plays the major role in bringing plasmatocytes to locations where they play essential developmental roles in the embryo, during larval life adhesion predominates and must be dynamically regulated to permit both normal proliferation and infectious responses. Plasmatocyte binding to tumors can lead to their inhibition or promote their invasion, depending on the genetic state of the tumor. In many of these steps similarities are evident to vertebrate macrophages and monocytes. The molecular mechanisms governing the movements, adhesion, and functions of the *Drosophila* immune system likely represent ancient programs upon which evolution has elaborated to permit the complex repertoire of immune cell behavior seen in vertebrates. Identifying new aspects of these mechanisms and their relevance for vertebrate immunology will occupy many exciting years ahead.

Acknowledgements

We thank all members of the Siekhaus lab for helpful discussions. We thank Matt DeGennaro, Matt Doerfler, Prashanth Rangan and Brian Stramer for comments on the manuscript. We apologize to all researchers whose work we did not have space to discuss. Daria Siekhaus is supported by Marie Curie CIG 34077/IRTIM and Aparna Ratheesh by Marie Curie IIF GA-2012-32950 BB: DICJL.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Luster AD, Alon R, Andrian von UH: **Immune cell migration in inflammation: present and future therapeutic targets.** *Nat Immunol* 2005, **6**:1182-1190.
 2. Munoz MA, Biro M, Weninger W: **T cell migration in intact lymph nodes in vivo.** *Curr Opin Cell Biol* 2014, **30**:17-24.
 3. Lemaitre B, Hoffmann J: **The host defense of *Drosophila melanogaster*.** *Annu Rev Immunol* 2007, **25**:697-743.
 4. Bunt S, Hooley C, Hu N, Scahill C, Weavers H, Skaer H: **Hemocyte-secreted type IV collagen enhances BMP signaling to guide renal tubule morphogenesis in *Drosophila*.** *Dev Cell* 2010, **19**:296-306.
 5. Sears HC, Kennedy CJ, Garrity PA: **Macrophage-mediated corpse engulfment is required for normal *Drosophila* CNS morphogenesis.** *Development* 2003, **130**:3557-3565.
 6. Zhou L, Hashimi H, Schwartz LM, Nambu JR: **Programmed cell death in the *Drosophila* central nervous system midline.** *Curr Biol* 1995, **5**:784-790.
 7. Woodcock KJ, Kierdorf K, Pouchelon CA, Vivancos V, Dionne MS, Geissmann F: **Macrophage-derived upd3 cytokine causes impaired glucose homeostasis and reduced lifespan in *Drosophila* fed a lipid-rich diet.** *Immunity* 2014 <http://dx.doi.org/10.1016/j.immuni.2014.12.023>.
 8. Braun A, Hoffmann JA, Meister M: **Analysis of the *Drosophila* host defense in *domino* mutant larvae, which are devoid of hemocytes.** *Proc Natl Acad Sci USA* 1998, **95**:14337-14342.
 9. Vlisidou I, Wood W: ***Drosophila* blood cells and their role in immune responses.** *FEBS J* 2015 <http://dx.doi.org/10.1111/febs.13235>.
 10. Costa A, Jan E, Sarnow P, Schneider D: **The Imd pathway is involved in antiviral immune responses in *Drosophila*.** *PLoS ONE* 2009, **4**:e7436.
 11. Pastor-Pareja JC, Wu M, Xu T: **An innate immune response of blood cells to tumors and tissue damage in *Drosophila*.** *Dis Model Mech* 2008, **1**:144-54 (discussion 153).
 12. Parisi F, Stefanatos RK, Strathdee K, Yu Y, Vidal M: **Transformed epithelia trigger non-tissue-autonomous tumor suppressor response by adipocytes via activation of Toll and Eiger/TNF signaling.** *Cell Rep* 2014, **6**:855-867.
- This interesting work shows the potential of *Drosophila* tumor models to study long and short-range interactions between epithelial tumors and the immune system. In an elegant and thorough series of genetic and imaging experiments, Parisi and colleagues show that tumor associated plasmatocytes (TAP) induce a feedback loop leading to tumor inhibition. Plasmatocytes produce the *Drosophila* TNF α ortholog, Eiger, which stimulates tumor production of a factor, Pvf1, that promotes plasmatocyte proliferation. Plasmatocytes also kill tumor cells by producing Eiger and the Toll ligand Spaetzle which induces Toll activation in the distant fat body; this analog of the mammalian liver produces some factor that acts with Eiger to facilitate tumor cell death.
13. Cho NK, Keyes L, Johnson E, Heller J, Ryner L, Karim F, Krasnow MA: **Developmental control of blood cell migration by the *Drosophila* VEGF pathway.** *Cell* 2002, **108**:865-876.
 14. Moreira CGA, Regan JC, Zaidman-Rémy A, Jacinto A, Prag S: ***Drosophila* hemocyte migration: an in vivo assay for directional cell migration.** *Methods Mol Biol* 2011, **769**:249-260.
 15. Stramer B, Wood W, Galko MJ, Redd MJ, Jacinto A, Parkhurst SM, Martin P: **Live imaging of wound inflammation in *Drosophila* embryos reveals key roles for small GTPases during in vivo cell migration.** *J Cell Biol* 2005, **168**:567-573.
 16. Wood W, Faria C, Jacinto A: **Distinct mechanisms regulate hemocyte chemotaxis during development and wound healing in *Drosophila melanogaster*.** *J Cell Biol* 2006, **173**:405-416.
 17. Makhijani K, Alexander B, Tanaka T, Rulifson E, Brückner K: **The peripheral nervous system supports blood cell homing and survival in the *Drosophila* larva.** *Development* 2011, **138**:5379-5391.
- In vertebrates the survival, proliferation and differentiation of hematopoietic cells are influenced by microenvironments including the peripheral nervous system (PNS). In this exciting paper, Makhijani and colleagues show for the first time a role for the PNS as a *Drosophila* hematopoietic niche supporting hemocyte homing and survival, indicating the potential of *Drosophila* to examine interactions of the nervous system and hematopoiesis.
18. Sampson CJ, Williams MJ: **Protocol for ex vivo incubation of *Drosophila* primary post-embryonic haemocytes for real-time analyses.** *Methods Mol Biol* 2012, **827**:359-367.
 19. Evans IR, Wood W: ***Drosophila* blood cell chemotaxis.** *Curr Opin Cell Biol* 2014, **30**:1-8.
 20. Holz A, Bossinger B, Strasser T, Janning W, Klapper R: **The two origins of hemocytes in *Drosophila*.** *Development* 2003, **130**:4955-4962.
 21. Lebestky T, Chang T, Hartenstein V, Banerjee U: **Specification of *Drosophila* hematopoietic lineage by conserved transcription factors.** *Science* 2000, **288**:146-149.
 22. de Velasco B, Mandal L, Mkrtychyan M, Hartenstein V: **Subdivision and developmental fate of the head mesoderm in *Drosophila melanogaster*.** *Dev Genes Evol* 2006, **216**:39-51.

23. Brückner K, Kockel L, Duchek P, Luque CM, Rørth P, Perrimon N: **The PDGF/VEGF receptor controls blood cell survival in *Drosophila***. *Dev Cell* 2004, **7**:73-84.
24. Siekhaus D, Haesemeyer M, Möffitt O, Lehmann R: **RhoL controls invasion and Rap1 localization during immune cell transmigration in *Drosophila***. *Nat Cell Biol* 2010, **12**:605-610.
This paper demonstrates that plasmotocytes move between epithelial cells when entering the germband in a genetically separable step of migration. This movement requires α -Integrin and a GEF for the Rap1 GTPase, raising potential parallels with vertebrate leukocyte vascular extravasation. RhoL was shown to be required for regulating Rap1 localization during this process.
25. Olofsson B, Page DT: **Condensation of the central nervous system in embryonic *Drosophila* is inhibited by blocking hemocyte migration or neural activity**. *Dev Biol* 2005, **279**:233-243.
26. Davies LC, Taylor PR: **Tissue-resident macrophages: then and now**. *Immunology* 2015, **144**:541-548.
27. Gentek R, Molawi K, Sieweke MH: **Tissue macrophage identity and self-renewal**. *Immunol Rev* 2014, **262**:56-73.
28. Godin I, Cumano A: **The hare and the tortoise: an embryonic haematopoietic race**. *Nat Rev Immunol* 2002, **2**:593-604.
29. Herbomel P, Thisse B, Thisse C: **Ontogeny and behaviour of early macrophages in the zebrafish embryo**. *Development* 1999, **126**:3735-3745.
30. Herbomel P, Thisse B, Thisse C: **Zebrafish early macrophages colonize cephalic mesenchyme and developing brain, retina, and epidermis through a M-CSF receptor-dependent invasive process**. *Dev Biol* 2001, **238**:274-288.
31. van Ham TJ, Kokel D, Peterson RT: **Apoptotic cells are cleared by directional migration and elmo1-dependent macrophage engulfment**. *Curr Biol* 2012, **22**:830-836.
32. Padrón-Barthe L, Temiño S, Villa del Campo C, Carramolino L, Isern J, Torres M: **Clonal analysis identifies homogenic endothelium as the source of the blood-endothelial common lineage in the mouse embryo**. *Blood* 2014, **124**:2523-2532.
33. Schulz C, Gomez Perdiguero E, Chorro L, Szabo-Rogers H, Cagnard N, Kierdorf K, Prinz M, Wu B, Jacobsen SEW, Pollard JW *et al.*: **A lineage of myeloid cells independent of Myb and hematopoietic stem cells**. *Science* 2012, **336**:86-90.
34. Ginhoux F, Greter M, Leboeuf M, Nandi S, See P, Gokhan S, Mehler MF, Conway SJ, Ng LG, Stanley ER *et al.*: **Fate mapping analysis reveals that adult microglia derive from primitive macrophages**. *Science* 2010, **330**:841-845.
35. Cuadros MA, Martin C, Coltey P, Almendros A, Navascués J: **First appearance, distribution, and origin of macrophages in the early development of the avian central nervous system**. *J Comp Neurol* 1993, **330**:113-129.
36. Rae F, Woods K, Sasmono T, Campanale N, Taylor D, Ovchinnikov DA, Grimmond SM, Hume DA, Ricardo SD, Little MH: **Characterisation and trophic functions of murine embryonic macrophages based upon the use of a Csf1r-EGFP transgene reporter**. *Dev Biol* 2007, **308**:232-246.
37. Gouon-Evans V, Rothenberg ME, Pollard JW: **Postnatal mammary gland development requires macrophages and eosinophils**. *Development (Cambridge, England)* 2000, **127**:2269-2282.
38. Watson CJ, Khaled WT: **Mammary development in the embryo and adult: a journey of morphogenesis and commitment**. *Development (Cambridge, England)* 2008, **135**:995-1003.
39. Ingman WV, Wyckoff J, Gouon-Evans V, Condeelis J, Pollard JW: **Macrophages promote collagen fibrillogenesis around terminal end buds of the developing mammary gland**. *Dev Dyn* 2006, **235**:3222-3229.
40. Davies LC, Jenkins SJ, Allen JE, Taylor PR: **Tissue-resident macrophages**. *Nat Immunol* 2013, **14**:986-995.
41. Parsons B, Foley E: **The *Drosophila* platelet-derived growth factor and vascular endothelial growth factor-receptor related (Pvr) protein ligands Pvf2 and Pvf3 control hemocyte viability and invasive migration**. *J Biol Chem* 2013, **288**:20173-20183.
42. Duchek P, Somogyi K, Jékely G, Beccari S, Rørth P: **Guidance of cell migration by the *Drosophila* PDGF/VEGF receptor**. *Cell* 2001, **107**:17-26.
43. Wang X, He L, Wu YI, Hahn KM, Montell DJ: **Light-mediated activation reveals a key role for Rac in collective guidance of cell movement in vivo**. *Nat Cell Biol* 2010, **12**:591-597.
44. Sopko R, Lin YB, Makhijani K, Alexander B, Perrimon N, Brückner K: **A systems-level interrogation identifies regulators of *Drosophila* blood cell number and survival**. *PLoS Genet* 2015, **11**:e1005056.
45. Tran TA, Kinch L, Peña-Llopis S, Kockel L, Grishin N, Jiang H, Brugarolas J: **Platelet-derived growth factor/vascular endothelial growth factor receptor inactivation by sunitinib results in Tsc1/Tsc2-dependent inhibition of TORC1**. *Mol Cell Biol* 2013, **33**:3762-3779.
46. McDonald JA, Pinheiro EM, Montell DJ: **PVF1, a PDGF/VEGF homolog, is sufficient to guide border cells and interacts genetically with Taiman**. *Development* 2003, **130**:3469-3478.
47. Donà E, Barry JD, Valentin G, Quirin C, Khmelinskii A, Kunze A, Durdu S, Newton LR, Fernández-Miñán A, Huber W *et al.*: **Directional tissue migration through a self-generated chemokine gradient**. *Nature* 2013, **503**:285-289.
48. Venkiteswaran G, Lewellis SW, Wang J, Reynolds E, Nicholson C, Knaut H: **Generation and dynamics of an endogenous, self-generated signaling gradient across a migrating tissue**. *Cell* 2013, **155**:674-687.
49. Boldajipour B, Mahabaleshwar H, Kardash E, Reichman-Fried M, Blaser H, Minina S, Wilson D, Xu Q, Raz E: **Control of chemokine-guided cell migration by ligand sequestration**. *Cell* 2008, **132**:463-473.
50. Shalaby F, Rossant J, Yamaguchi TP, Gertsenstein M, Wu X-F, Breitman ML, Schuh AC: **Failure of blood-island formation and vasculogenesis in Flk-1-deficient mice**. *Nature* 1995, **376**:62-66.
51. Hidaka M, Stanford WL, Bernstein A: **Conditional requirement for the Flk-1 receptor in the in vitro generation of early hematopoietic cells**. *Proc Natl Acad Sci USA* 1999, **96**:7370-7375.
52. Kaminski WE, Lindahl P, Lin NL, Broudy VC, Crosby JR, Hellström M, Swolin B, Bowen-Pope DF, Martin PJ, Ross R *et al.*: **Basis of hematopoietic defects in platelet-derived growth factor (PDGF)-B and PDGF β -receptor null mice**. *Blood* 2001, **97**:1990-1998.
53. Clauss M, Gerlach M, Gerlach H, Brett J, Wang F, Familletti PC, Pan YC, Olander JV, Connolly DT, Stern D: **Vascular permeability factor: a tumor-derived polypeptide that induces endothelial cell and monocyte procoagulant activity, and promotes monocyte migration**. *J Exp Med* 1990, **172**:1535-1545.
54. Clauss M, Weich H, Breier G, Knies U, Röckl W, Waltenberger J, Risau W: **The vascular endothelial growth factor receptor Flt-1 mediates biological activities. Implications for a functional role of placenta growth factor in monocyte activation and chemotaxis**. *J Biol Chem* 1996, **271**:17629-17634.
55. Deuel TF, Senior RM, Huang JS, Griffin GL: **Chemotaxis of monocytes and neutrophils to platelet-derived growth factor**. *J Clin Invest* 1982, **69**:1046-1049.
56. Krettek A, Ostergren-Lundén G, Fager G, Rosmond C, Bondjers G, Lustig F: **Expression of PDGF receptors and ligand-induced migration of partially differentiated human monocyte-derived macrophages. Influence of IFN- γ and TGF- β** . *Atherosclerosis* 2001, **156**:267-275.
57. Wang JM, Collela S, Allavena P, Mantovani A: **Chemotactic activity of human recombinant granulocyte-macrophage colony-stimulating factor**. *Immunology* 1987, **60**:439.
58. Pixley FJ: **Macrophage migration and its regulation by CSF-1**. *Int J Cell Biol* 2012:501962.

59. Huelsmann S, Hepper C, Marchese D, Knöll C, Reuter R: **The PDZ-GEF Dizzy regulates cell shape of migrating macrophages via Rap1 and integrins in the *Drosophila* embryo.** *Development* 2006, **133**:2915-2924.
60. Abram CL, Abram CL, Lowell CA, Lowell CA: **The ins and outs of leukocyte integrin signaling.** *Annu Rev Immunol* 2009, **27**:339-362.
61. Nourshargh S, Hordijk PL, Sixt M: **Breaching multiple barriers: leukocyte motility through venular walls and the interstitium.** *Nat Rev Mol Cell Biol* 2010, **11**:366-378.
62. Stramer B, Moreira S, Millard T, Evans I, Huang C-Y, Sabet O, Milner M, Dunn G, Martin P, Wood W: **Clasp-mediated microtubule bundling regulates persistent motility and contact repulsion in *Drosophila* macrophages in vivo.** *J Cell Biol* 2010, **189**:681-689.
- Using beautiful live imaging and analysis Stramer and colleagues show for the first time a role for aligned microtubule arms in mediating cell-cell repulsion during polarized cell migration *in vivo* and demonstrate that this repulsion is essential for the proper dispersal of plasmatocytes in later embryonic stages. This work highlights the importance of studying microtubule dynamics *in vivo* since primary hemocytes plated *in vitro* do not form these microtubule arms.
63. Davis JR, Luchici A, Mosis F, Thackery J, Salazar JA, Mao Y, Dunn GA, Betz T, Miodownik M, Stramer BM: **Inter-cellular forces orchestrate contact inhibition of locomotion.** *Cell* 2015 <http://dx.doi.org/10.1016/j.cell.2015.02.015>.
- This paper shows the power of sophisticated live imaging and analysis in dissecting the cell biological changes that underlie plasmatocyte behaviors. Davis *et al.* investigate the mechanisms underlying the movement of two plasmatocytes away from one another after contact. They demonstrate that formation of a zyxin containing adhesion precedes a localized reduction in actin flow, formation of an actin stress fiber and aligned bundles of microtubules. This leads to a build up of tension between the coupled cells; its release appears to underlie the swift movement of the cells away from one another.
64. Comber K, Huelsmann S, Evans I, Sánchez-Sánchez BJ, Chalmers A, Reuter R, Wood W, Martin-Bermudo MD: **A dual role for the β PS integrin *myospheroid* in mediating *Drosophila* embryonic macrophage migration.** *J Cell Sci* 2013, **126**:3475-3484.
65. Welman A, Serrels A, Brunton VG, Ditzel M, Frame MC: **Two-color photoactivatable probe for selective tracking of proteins and cells.** *J Biol Chem* 2010, **285**:11607-11616.
66. Babcock DT, Brock AR, Fish GS, Wang Y, Perrin L, Krasnow MA, Galko MJ: **Circulating blood cells function as a surveillance system for damaged tissue in *Drosophila* larvae.** *Proc Natl Acad Sci USA* 2008, **105**:10017-10022.
67. Lanot R, Zachary D, Holder F, Meister M: **Postembryonic hematopoiesis in *Drosophila*.** *Dev Biol* 2001, **230**:243-257.
68. Bretscher AJ, Honti V, Binggeli O, Burri O, Poidevin M, Kurucz É, Zsámboki J, Andó I, Lemaitre B: **The Nimrod transmembrane receptor Eater is required for hemocyte attachment to the sessile compartment in *Drosophila melanogaster*.** *Biol Open* 2015 <http://dx.doi.org/10.1242/bio.201410595>.
- This work contains the surprising and intriguing finding that the transmembrane protein Eater plays two divergent roles in plasmatocytes. It is required both for phagocytosis of gram positive bacteria and binding to the hematopoietic pockets on the larval body wall where plasmatocytes become sessile and proliferate. Bretscher *et al.* show that Eater is required in plasmatocytes themselves both for their own localization and that of crystal cells.
69. Leitão AB, Sucena É: ***Drosophila* sessile hemocyte clusters are true hematopoietic tissues that regulate larval blood cell differentiation.** *Elife* 2015:4.
70. Galko MJ, Krasnow MA: **Cellular and genetic analysis of wound healing in *Drosophila* larvae.** *PLoS Biol* 2004, **2**:E239.
71. Sorrentino RP, Carton Y, Govind S: **Cellular immune response to parasite infection in the *Drosophila* lymph gland is developmentally regulated.** *Dev Biol* 2002, **243**:65-80.
72. Márkus R, Laurinyecz B, Kurucz E, Honti V, Bajusz I, Sipos B, Somogyi K, Kronhamn J, Hultmark D, Andó I: **Sessile hemocytes as a hematopoietic compartment in *Drosophila melanogaster*.** *Proc Natl Acad Sci USA* 2009, **106**:4805-4809.
73. Schmid MR, Anderl I, Vesala L, Vanha-Aho L-M, Deng X-J, Rämét M, Hultmark D: **Control of *Drosophila* blood cell activation via Toll signaling in the fat body.** *PLoS One* 2014, **9**:e102568.
74. Zettervall C-J, Anderl I, Williams MJ, Palmer R, Kurucz É, Andó I, Hultmark D: **A directed screen for genes involved in *Drosophila* blood cell activation.** *Proc Natl Acad Sci USA* 2004, **101**:14192-14197.
75. Kocks C, Cho JH, Nehme N, Ulvila J, Pearson AM, Meister M, Strom C, Conto SL, Hetru C, Stuart LM *et al.*: **Eater, a transmembrane protein mediating phagocytosis of bacterial pathogens in *Drosophila*.** *Cell* 2005, **123**:335-346.
76. Chung Y-SA, Kocks C: **Recognition of pathogenic microbes by the *Drosophila* phagocytic pattern recognition receptor Eater.** *J Biol Chem* 2011, **286**:26524-26532.
77. Trombetta ES, Mellman I: **Cell biology of antigen processing in vitro and in vivo.** *Annu Rev Immunol* 2005, **23**:975-1028.
78. Makhijani K, Brückner K: **Of blood cells and the nervous system: hematopoiesis in the *Drosophila* larva.** *Fly (Austin)* 2012, **6**:254-260.
79. Gomez Perdiguero E, Klapproth K, Schulz C, Busch K, Azzoni E, Crozet L, Garner H, Trouillet C, de Bruijn MF, Geissmann F *et al.*: **Tissue-resident macrophages originate from yolk-sac-derived erythro-myeloid progenitors.** *Nature* 2015, **518**:547-551.
80. Hauling T, Krautz R, Márkus R, Volkenhoff A, Kucerova L, Theopold U: **A *Drosophila* immune response against Ras-induced overgrowth.** *Biol Open* 2014, **3**:250-260.
81. Bilder D: **Epithelial polarity and proliferation control: links from the *Drosophila* neoplastic tumor suppressors.** *Genes Dev* 2004, **18**:1909-1925.
82. Pagliarini RA, Xu T: **A genetic screen in *Drosophila* for metastatic behavior.** *Science* 2003, **302**:1227-1231.
83. Cordero JB, Macagno JP, Stefanatos RK, Strathdee KE, Cagan RL, Vidal M: **Oncogenic Ras diverts a host TNF tumor suppressor activity into tumor promoter.** *Dev Cell* 2010, **18**:999-1011.
- Work by Cordero and colleagues reveals intriguing similarities in the functions of the *Drosophila* TNF α homolog, Eiger, to its vertebrate counterpart in tumor promotion. Eiger mediated JNK signaling induced by tumor associated plasmatocytes has previously been shown to result in tumor cell death and inhibition. Cordero *et al* demonstrate for the first time that over expression of oncogenic Ras results in the hijacking of this pathway to convert an *in situ* tumor into an invasive one.
84. Ostuni R, Kratochvill F, Murray PJ, Natoli G: **Macrophages and cancer: from mechanisms to therapeutic implications.** *Trends Immunol* 2015 <http://dx.doi.org/10.1016/j.it.2015.02.004>.